

TUBERCULOSIS AND HIV

8 countries account for **70%** of all TB deaths among people living with HIV

ANNUAL GLOBAL FUNDING FOR TUBERCULOSIS IS ALMOST **US\$ 3 BILLION** SHORT OF WHAT IS REQUIRED

IN 2015, 10.4 MILLION PEOPLE FELL ILL WITH TB AND 1.8 MILLION DIED FROM THE DISEASE



TB IS THE **LEADING CAUSE OF DEATH** AMONG PEOPLE LIVING WITH HIV

57% OF HIV-ASSOCIATED TB CASES WENT **UNTREATED** IN 2015



In 2015, around 400 000 people died from AIDS-related TB, including **40 000 children**

UNAIDS IS WORKING WITH PARTNERS TO REDUCE TB-ASSOCIATED DEATHS AMONG PEOPLE LIVING WITH HIV **BY 75% BY 2020**

SIMPLE, AFFORDABLE AND EFFECTIVE HIV/TB PROGRAMMES

All people living with HIV should have access to:

- Antiretroviral Therapy
- TB diagnostics and treatment
- Regular TB screening
- TB preventive therapy (if no TB symptoms)

All people living with TB should have access to:

- HIV testing and antiretroviral therapy
- HIV prevention options
- TB treatment

