

TUBERCULOSIS AND HIV

IN 2017, 10 MILLION PEOPLE FELL ILL WITH TB AND 1.6 MILLION DIED FROM THE DISEASE



People living with HIV are up to **20 times** more likely to fall ill **with TB**

ANNUAL GLOBAL FUNDING FOR TUBERCULOSIS IS **US\$ 3.5 BILLION** SHORT OF WHAT IS REQUIRED



TB IS THE **LEADING CAUSE OF DEATH** AMONG PEOPLE LIVING WITH HIV

UNAIDS IS WORKING WITH PARTNERS TO REDUCE TB-ASSOCIATED DEATHS AMONG PEOPLE LIVING WITH HIV **BY 75% BY 2020**



In 2017, approximately **300 000** people died from AIDS-related TB

TB IS CURABLE:
54 MILLION
LIVES HAVE BEEN SAVED **SINCE 2000**

SIMPLE, AFFORDABLE AND EFFECTIVE HIV/TB PROGRAMMES

All people living with HIV should have access to:

- Antiretroviral Therapy
- TB diagnostics and treatment
- Regular TB screening
- TB preventive therapy (if no TB symptoms)



All people living with TB should have access to:

- HIV testing and antiretroviral therapy
- TB treatment
- HIV prevention options