





TRANSGENDER WOMEN WHO HAVE SEX WITH MALES PHILIPPINES


WHO WE SURVEYED




Median age (range)
23
(15-65)



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



FEMINIZING ENHANCEMENTS



RISK BEHAVIORS IN THE PAST 12 MONTHS



INCLUSION CRITERIA
Assigned sex at birth is male, 15 years or older, who reported oral or anal sex with a male in the past 12 months

IMPORTANT NOTES
Data presented is among those who identify themselves as female, a subset of the 2018 IHSS for MSM & TGW
Data presented is the aggregate of the 13 cities: Angeles, Baguio, Cagayan de Oro, Davao, Gensan, Iloilo, Zamboanga, Talisay, Pasay, Quezon City, Taguig
Time location sampling was used
Data presented is adjusted using sample weights
*Tested in the past 12 months and know their status also include those who know they are HIV-positive even if their last test was more than 12 months ago
HIV prevalence is among those who ever had anal sex



TRANSGENDER WOMEN WHO HAVE SEX WITH MALES PHILIPPINES

KNOWLEDGE ON HIV

33%

Know all the five basic facts of HIV transmission and prevention



84%
Know that a healthy looking person can have HIV



66%
Know that a person cannot get HIV from mosquito bites



81%
Know that using condoms reduce the risk of HIV transmission



77%
Know that a person cannot get HIV from using toilet bowls or urinals in public places



68%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



Aware that there is HIV treatment




Aware that there is a local Social Hygiene Clinic (SHC)



Knows that sharing needles used by an HIV infected person can increase the risks of HIV transmission

CONDOM USE AND ACCESS



● No Condom Access ● Buys Condoms
● Gets Free Condoms ● Gets Free and Buys Condoms

39%
Condom use during last anal sex (past 12 months)

15%
Consistent condom use (last 3 anal sex partners in the past 12 months)

REASONS FOR NOT USING CONDOMS DURING LAST ANAL SEX

Unplanned sex	37%
No sensation	28%
Partner did not want	21%
Don't want to carry around	11%
Trusted partner is negative	7%
Caught in the moment	7%
Embarrassed to buy	6%
Only one partner	6%
Minor/underage	3%
Too expensive	3%
Don't know where to get	2%
Don't know where to buy	2%
Don't know how to use	1%

HIV TESTING AND STI PREVALENCE



57%




42%




41%


Syphilis



Hepatitis B



HIV



*95% confidence interval

INCLUSION CRITERIA

Assigned sex at birth is male, 15 years or older, who reported oral or anal sex with a male in the past 12 months

IMPORTANT NOTES

Data presented is among those who identify themselves as female, a subset of the 2018 IHBS for MSM & TGW
Data presented is the aggregate of the 13 cities: Angeles, Baguio, Cagayan de Oro, Davao, Gensan, Iloilo, Zamboanga, Talisay, Pasay, Quezon City, Taguig
Time location sampling was used
Data presented is adjusted using sample weights
*Tested in the past 12 months and know their status also include those who know they are HIV- positive even if their last test was more than 12 months ago
HIV prevalence is among those who ever had anal sex



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines
Tel: +632 651-7800 loc 2952
Email: hivepicenter@gmail.com
Website: www.doh.gov.ph

