Bangladesh

Tuberculosis profile

Population 2017 165 million

		Rate
Estimates of TB burden*, 2017	Number (thousands)	(per 100 000 population)
Mortality (excludes HIV+TB)	59 (38–85)	36 (23–52)
Mortality (HIV+TB only)	0.17 (0.085–0.29)	0.11 (0.05–0.18)
Incidence (includes HIV+TB)	364 (265-479)	221 (161–291)
Incidence (HIV+TB only)	0.55 (0.27-0.92)	0.33 (0.17-0.56)
Incidence (MDR/RR-TB)**	8.4 (3.8–15)	5.1 (2.3–9)

Estimated TB incidence by age and sex (thousands)*, 2017			
	0-14 years	> 14 years	Total
Females	17 (16–18)	118 (98–137)	134 (110–158)
Males	18 (17–19)	212 (164–259)	230 (176–284)
Total	35 (32–38)	329 (237-421)	364 (265-479)

TB case notifications, 2017	
Total cases notified	244 201
Total new and relapse	242 639
- % tested with rapid diagnostics at time of diagnosis	<1%
- % with known HIV status	2%
- % pulmonary	81%
- % bacteriologically confirmed among pulmonary	74%
Universal health coverage and social protection	
TB treatment coverage (notified/estimated incidence), 2017	67% (51–92)

TB case fatality ratio (estimated mortality/estimated incidence), 2017	0.17 (0.1	0.17 (0.1–0.26)	
TB/HIV care in new and relapse TB patients, 2017	Number	(%)	
Patients with known HIV-status who are HIV-positive	89	2%	
- on antiretroviral therapy	84	94%	

		Dunisias sales dun nto al	Tatal
		Previously treated	Total
Drug-resistant TB care, 2017	New cases	cases	number***
Estimated MDR/RR-TB cases among notified			5 800
pulmonary TB cases		(3	800–7800)
Estimated % of TB cases with MDR/RR-TB	1.6% (0.74-2.8)	29% (24–35)	
% notified tested for rifampicin resistance	18%	63%	49 943
MDR/RR-TB cases tested for resistance to second	ond-line drugs		362
Laboratory-confirmed cases		MDR/RR-TB: 944,	XDR-TB: 6
Patients started on treatment ****		MDR/RR-TB: 920,	XDR-TB: 6

Treatment success rate and cohort size	Success	Cohort
New and relapse cases registered in 2016	94%	222 252
Previously treated cases, excluding relapse, registered in 2016	86%	1 669
HIV-positive TB cases registered in 2016	49%	87
MDR/RR-TB cases started on second-line treatment in 2015	78%	880
XDR-TB cases started on second-line treatment in 2015		0

MDR/RR-1B cases started on second-line treatment in 2015	78%	880
XDR-TB cases started on second-line treatment in 2015		0
TB preventive treatment, 2017		
% of HIV-positive people (newly enrolled in care) on preventive treatment		
% of children (aged < 5) household contacts of bacteriologically-confirmed	21%	(19–23)

TB financing, 2018	
National TB budget (US\$ millions)	66
Funding source: 14% domestic, 55% international, 31% unfunded	

* Ranges represent uncertainty intervals

TB cases on preventive treatment

TB patients facing catastrophic total costs

- ** MDR is TB resistant to rifampicin and isoniazid; RR is TB resistant to rifampicin
- *** Includes cases with unknown previous TB treatment history
- **** Includes patients diagnosed before 2017 and patients who were not laboratory-confirmed









