

Survey: Behaviorual Surveillance Among Drug Injectors in Indonesia, 2002

Title:	Behaviorual Surveillance Among Drug Injectors in Indonesia, 2002													
Survey Type:	BSS													
Country:	Indonesia													
Year:	2002													
Start Date:	[n/a]													
End Date:	[n/a]													
Respondents:	<table border="1"> <thead> <tr> <th>Female</th> <th>Male</th> </tr> </thead> <tbody> <tr> <td>Respondent: [n/a]</td> <td>Respondent: Target Groups</td> </tr> <tr> <td>Age: [n/a]</td> <td>Age: 18 +</td> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: 650</td> </tr> <tr> <th>Facilities</th> <th>Households</th> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: [n/a]</td> </tr> </tbody> </table>	Female	Male	Respondent: [n/a]	Respondent: Target Groups	Age: [n/a]	Age: 18 +	Sample Size: [n/a]	Sample Size: 650	Facilities	Households	Sample Size: [n/a]	Sample Size: [n/a]	
Female	Male													
Respondent: [n/a]	Respondent: Target Groups													
Age: [n/a]	Age: 18 +													
Sample Size: [n/a]	Sample Size: 650													
Facilities	Households													
Sample Size: [n/a]	Sample Size: [n/a]													
Implementing:	Indonesia Ministry of Health, University of Indonesia, FHI													
Description:	BSS among men aged 18 years or more who have injected drugs in the last month. Survey was conducted in three large cities: Jakarta, Surabaya, and Bandung. Questions asked cover areas such as: Sexual Activity, Condom Use, Injecting Drug Use, HIV Testing.													
Footnote:	[n/a]													

RESULTS

Indicators:	<p>4.0.1) Heard of HIV/AIDS 99%</p> <p>4.1.2) Knowledge of HIV prevention methods - Use of condoms (prompted) 91%</p> <p>4.2.1) No incorrect beliefs about AIDS - Healthy-looking person can have the AIDS virus 11%</p> <p>5.1) Population requesting an HIV test, receiving a test and receiving test results 13%</p> <p>5.1.1) Population receiving an HIV test 22%</p> <p>8.1) Higher risk sex in the last year 42% , n=444</p> <p>8.2) Condom use at last higher risk sex 17%, n=189</p> <p>8.2.1) Condom use at last sex with a spouse or cohabiting partner 14%, n=153</p> <p>8.3) Commercial sex in last year =40%</p> <p>8.4) Condom use at last commercial sex, reported by client 23%, n=263</p> <p>8.7) Condom use at last anal sex between men</p> <p>10.1) Injecting drug users sharing equipment at last injection 62%</p>
--------------------	---