Cambodia HIV Country Profile 2022


WHO/UCN/HHS/21.10


Year


4.0%


Value

Cambodia











Estimated number of people dying from HIV -related causes





Trend since 2010 Epidemiological HIV data (2021)

Estimated number of women (15+) living with HIV 39 000 Estimated number of people newly infected with HIV 1 100 Estimated number of people living with HIV 74 000 Estimated number of people dying from HIV-related causes 1 100 Estimated number of children aged 0 to 14 living with HIV 2 300 Estimated incidence rate per 1000 uninfected population 0.07 Estimated adult (15-49) prevalence 0.6% Elimination of mother-to-child transmission (EMTCT) of HIV &

Trend since 2010 (-36%)

2 000

Value

Reported number of pregnant women living with HIV who received antiretrovirals for PMTCT	472	202
Final transmission rate including breastfeeding period ³ (%)	13.6%	202
Estimated percentage of pregnant women living with HIV who received antiretrovirals for PMTCT ² (%)	80%	202
Antenatal care coverage - at least one visit (%)	95.3%	201
Antenatal care coverage - at least four visits (%)	75.6%	201
Antenatal care attendees who were tested for syphilis (%)	79.7%	201
Antenatal care attendees who tested positive for syphilis and who received benzathine penicillin IM (%)	84.5%	201
Antenatal care attendees who tested positive for syphilis (%)	0.1%	201
% of pregnant women with known HIV status	95%	202
% of infants born to women living with HIV receiving a virological test within two months of birth (EID) ¹ (%)	13.7%	202

84% Estimated ART coverage Estimated ART coverage (adults 15+) 85% Estimated ART coverage (children, 0-14)¹ 56% Reported number of children receiving ART 1 265 Reported number of people living with HIV receiving ART 62 561 Reported number of people living with HIV receiving ART (adults 15+) 61 296

HIV prevalence among men who have sex with men (%)


Antiretroviral therapy (ART) (2021)

targets: 190% 295% 3Less than 5 Key populations (2021)

Tracing indicators for elimination of mother-to-child transmission. A color scheme

(green, orange, red) is used to indicate if the country is on target to meet 2020

HIV prevalence among people who inject drugs (%)	15.2%		
HIV prevalence among sex workers (%)	N/A		
Needles distributed per person who inject drugs per year	168		
Percentage condom use among sex workers with most recent client	N/A		
Percentage of people who inject drugs receiving OST	6.4%		
Median annual price o	of ARV drug regimens for adults		
Wicalan annual price of All Variaging Teginnens			
First line	Second line		
◆[TDF+3TC+EFV] [600+300+300]mg ◆[ZDV+3TC+NVP] [150+200+300]mg	◆[TDF+3TC]+[ATV/r] ◆[TDF+3TC]+[LPV/r] ◆[TDF+FTC]+[LPV/r] ◆[ZDV+3TC]+[◆[ZDV+3TC]+[
L dear	ğ 400 ••••••••••••••••••••••••••••••••••		



National HIV policies and plans (2022)			
Indicator	Value Value		
Implementation of national policy on HIV self-testing (HIVST)	HIVST policy and implementation		
Treatment initiation threshold among adults and adolescents	Treat all regardless of CD4 count		
Implementation of treat all policy among adults and adolescents	Implemented countrywide (>95% of treatment sites)		
Implementation of lifelong ART for pregnant & breastfeeding women	Implemented		
Treatment initiation threshold among children	Treat all, regardless of age		
Adoption of WHO 2017 recommendation on rapid initiation of ART	Rapid initiation within 7 days of HIV diagnosis		
National policy supporting community delivery of ART	Policy implemented at pilot sites		
Adoption of WHO recommendation to offer package of interventions for patients with advanced HIV disease	Fully adopted •		
Policy on the frequency of clinic visits for adults who are doing well on antiretroviral therapy	Every 3 months for adults who are doing well on ART		
Policy on how frequently adults who are doing well on antiretroviral therapy should pick-up antiretroviral medicine (MMD)	Every 3 months for adults who are doing well on ART		
Dolutegravir (DTG) as part of 1st line ARVs for adults and adolescents	DTG introduced in national guidelines and procurement initiated		
Implementation of national policy on viral load monitoring	Implemented countrywide (>95% of treatment sites)		
Availability of point-of-care viral load testing	Not available		
National M&E plan or strategy for HIV	Integrated HIV monitoring and evaluation strategy or plan		

Sources: Global AIDS Monitoring UNAIDS/WHO/UNICEF (2022), Global Health Expenditure Database WHO (2022), Global Health Observatory WHO (2022), Global Price Reporting Mechanism WHO (2022), Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat (2022), UNICEF Data (2022), United Nations Development Programme (2022), Spectrum estimates UNAIDS/WHO (2022), World Bank (2022)

Definition of acronyms: PLHIV = People living with HIV, HIV = Human Immunodeficiency Virus, AIDS = Acquired Immunodeficiency Syndrome, ART = Antiretroviral Therapy,

ARVs = Antiretrovirals, PMTCT = Prevention of Mother-to-Child Transmission, EID = Early Infant Diagnosis, OST = Opioid Substitution Therapy, MMD = Multi-month Dispensing