



Newly Diagnosed HIV Cases in the Philippines

In February 2011, there were 159 new HIV Ab sero-positive individuals confirmed by the STD/AIDS Cooperative Central Laboratory (SACCL) and reported to the HIV and AIDS Registry (Table 1). This was a 22% increase compared to the same period last year (n=130 in 2010) [Figure 1].

Most of the cases (91%) were males. The median age was 28 years (age range:15-56 years). The 20-29 year (52%) age-group had the most number of cases. Fifty-six percent (89) of the reported cases were from the National Capital Region (NCR).

Reported mode of transmission was sexual contact (155) and blood transfusion (1). Three did not report mode of transmission [Table 2, page 3]. Males having sex with other Males (79%) were the predominant type of sexual transmission [Figure 2]. Most (98%) of the cases were still asymptomatic at the time of reporting [Figure 3].

AIDS Cases

Of the 159 HIV positive cases, three were reported as AIDS cases. Two were males and one was female. The median age was 24 years (age range: 22-27 years). All acquired infection through sexual contact (1 heterosexual, 1 homosexual, and 1 bisexual). There was one reported death for this month.

Overseas Filipino Workers (OFW)

Nineteen of the 159 (12%) reported cases were OFWs [Figure 9, page 3]. Fifteen were males and four were females. The median age was 32 years (age range: 25-41 years). All cases acquired the HIV infection through sexual contact (7 heterosexual, 6 homosexual, and 6 bisexual).

Table 1. Quick Facts

Demographic Data	Feb 2011	Jan-Feb 2011	Cumulative Data: 1984–2011
Total Reported Cases	159	311	6,326
Asymptomatic Cases	156	305	5,463
AIDS Cases	3	6	863
Males	144	285	4,984*
Females	15	26	1,331*
Youth 15-24yo	44	89	1,302
Children <15yo	0	0	55
Reported Deaths due to AIDS	1	1	324

*Note: No data available on sex for eleven (11) cases.

Figure 1. Number of New HIV Cases per Month (2008-2011)




Figure 2. Comparison of the Proportion of Types of Sexual Transmission in 2011, 2010 & Cumulative Data (1984-2011)





Figure 3. Number of HIV/AIDS Cases Reported in the Philippines by Year, Jan 1984 to February 2011 (N=6,326)




*Five initially asymptomatic cases reported in 2008, died due to AIDS that same year.

AIDS Cases (1984-2011)

Of the 311 HIV positive cases in 2011, six were reported as AIDS cases. Eighty-three percent were males. Ages ranged from 22-51 years (median 32 years). All acquired the infection through sexual contact [heterosexual (2), homosexual contact (3), and bisexual contact (1)].

From 1984 to 2011, there were 863 AIDS cases reported, 71% (615) were males. Median age was 35 years (range 1-72 years). Of the AIDS cases, there were 324 (38%) deaths during the reported period. Sexual contact was the most common mode of HIV transmission, accounting for 93% (802) of all AIDS cases. More than half (450) of sexual transmission was through heterosexual contact, followed by homosexual contact (275) then bisexual contact (77). Other modes of transmission include: mother-to-child transmission (17), blood transfusion (10), injecting drug use (4), and needle prick injuries (2) [Figure 4]. Three percent (28) of the AIDS cases did not report mode of HIV transmission.

Fig 4. Proportion of Modes of Transmission of AIDS Cases by Year, Jan 1984–February 2011



*Note: 28 did not report mode of transmission

Demographic Characteristics (1984-2011)

In 2011, there were a total of 311 cases reported. 92% of the cases reported were males (285). Ages ranged from 15-61 years old (median 28 years). The 20-29 year old age group (59%) had the most number of cases for 2011. For the male age group, the most number of cases were found among the 20-24 years old (27%), 25-29 years old (32%) and 30-34 years old (21%) [Figure 5].

From 1984 to 2011, there were 6,326 HIV Ab sero-positive cases reported (Table 1), of which 5,463 (86%) were asymptomatic and 863 (14%) were AIDS cases. As shown in Figure 6, there is a significant difference in the number of male and female cases reported. Seventy-nine percent (4,984) were males. Ages ranged from 1-73 years (median 30 years). The age groups with the most number of cases were: 20-24 years (19%), 25-29 (25%) and 30-34 years (19%) [Figure 6].

Figure 5. Proportion of Sex & Age-Groups in Feb 2011 & Jan-Feb 2011





Figure 6. Comparison of the Distribution of Male and Female HIV Cases by Age-Group and Certain Highlighted Years



Modes of Transmission (1984-2011)

In 2011, 98% (306) were infected through sexual contact and <1% (1) through blood transfusion; 1% (4) had no reported data on mode of transmission (Table 2). There were 281 males and 25 females infected through sexual transmission. Their ages ranged from 15-61 years old (median 28 years) [Figure 7].


Of the 6,326 with HIV from 1984 to 2011, 90% (5,712) were infected through sexual contact, 1% (52) through mother-to-child transmission and 3% (155) through needle sharing among injecting drug users. Other modes of transmission are listed in Table 2. No data is available for 6% (384) of the cases. Cumulative data shows 45% (2,546) were infected through heterosexual contact, 35% (1,993) through homosexual contact, and 21% (1,173) through bisexual contact. From 2007 there has been a shift in the predominant trend of sexual transmission from heterosexual contact (25%) to males having sex with other males (75%) [Figure 8].

Table 2. Reported Mode of HIV Transmission

Mode of Transmission	Feb 2011 n=159	Jan–Feb 2011 n=311	Cumulative N=6,326
Sexual Contact	155	306	5,712
<i>Heterosexual contact</i>	33 (21%)	61 (20%)	2,546 (45%)
<i>Homosexual contact</i>	70 (45%)	144 (47%)	1,993 (35%)
<i>Bisexual contact</i>	52 (34%)	101 (33%)	1,173 (21%)
Blood/Blood Products	1*	1	20
Injecting Drug Use	0	0	155
Needle Prick Injury	0	0	3
Mother-to-Child	0	0	52
No Data Available	3	4	384


*Details for verification

Figure 7. Proportion of Modes of HIV Transmission by Age-Group, 2011(n=311)



*No data available on Modes of Transmission for four (4) cases

Figure 8. Proportion of Types of Sexual Transmission, Jan 1984–Feb 2011



Overseas Filipino Workers (OFW)


In 2011, there were 36 HIV positive OFWs, comprising 12% of cases reported for the year [Figure 9]. Of these, 30 (83%) were males and 6 (17%) were females; all infected through sexual contact.

There were 1,558 HIV positive OFWs since 1984, comprising 25% of all reported cases [Figure 9]. Seventy-five percent (1,175) were males. Ages ranged from 18 to 69 years (median 36 years). Sexual contact (96%) was the predominant mode of transmission (Table 3). Eighty-three percent (1,291) were asymptomatic while 17% (267) were AIDS cases.

Table 3. Reported Mode of HIV Transmission Among OFWs

Mode of Transmission	Feb 2011 n= 19	Jan– Feb 2011 n= 36	Cumulative N=1,558
Sexual Transmission	19	36	1,496
<i>Heterosexual contact</i>	7 (37%)	13 (36%)	1,014 (68%)
<i>Homosexual contact</i>	6 (32%)	10 (28%)	288 (19%)
<i>Bisexual contact</i>	6 (32%)	13 (36%)	194 (13%)
Blood/Blood Products	0	0	10
Injecting Drug Use	0	0	1
Needle Prick Injury	0	0	3
No Data Available	0	0	48

Figure 9. Number of OFWs Compared to Non-OFWs by Year (1984-2011*)



*Data includes January to February 2011 only.

Program Related Information

Of the 159 HIV positive cases reported in February 2011, three were classified as AIDS. Seventy percent of the cases received information on HIV prevention, services available for HIV cases, implications of an HIV positive result from screening and confirmation. Their sources of information were one-on-one counseling, group counseling, pre-departure orientation seminar (PDOS), pamphlets, videos, internet and seminars.

Blood Units Screened for HIV

Note: The following information is from the National Voluntary Blood Safety Program (NVBSP) which monitors blood safety of donated blood. HIV reactive blood units are referred to the Research Institute for Tropical Medicine (RITM) for confirmation. RITM is the National Reference Laboratory for the NVBSP.

From January to February 2011, 26 blood units were confirmed to be positive by the RITM.

For February 2011, out of the 67 blood units referred for HIV confirmation, 15 units were positive for HIV and 50 units were negative for HIV; 2 units had indeterminate result [Table 4].

Figure 10. HIV Positive Blood Units by Month & Year (2008-2011)




Table 4. Results of Blood Units Referred for HIV Confirmation

Monthly Report	2011		
	Blood units* referred	Positive	Indeterminate
January	85	11	0
February	67	15	2
March			
April			
May			
June			
July			
August			
September			
October			
November			
December			
Total for the year (Jan –Feb only)	152	26	2

* One blood donor can donate more than one blood unit.

** These are HIV positive blood units, not donors. Donors of HIV positive blood units may or may not be in the HIV & AIDS Registry.



National HIV/AIDS & STI Strategic Information and Surveillance Unit

National Epidemiology Center,
Department of Health, Bldg. 9,
San Lazaro Compound,
Sta. Cruz, Manila 1003 Philippines

Tel: +632 743 8301 local 1900 to 1907
Fax: +632 743 6076 / 743 1937
Email: HIVepicenter@gmail.com
Website: http://www.doh.gov.ph

Philippine HIV & AIDS Registry Report Editorial Team:

Ma. Lourdes S. Macapanpan
Ma. Lourdes S. Macapanpan
HIV Surveillance Assistant, HIV Unit

Noel S. Pineda
Noel S. Pineda, RN
HIV Surveillance Officer, HIV Unit

Genesis Jay J. Samonte
Genesis Jay J. Samonte, MD, PHSAE
Epidemiologist, HIV Unit

Agnes B. Segarra
Agnes B. Segarra, MD, PHSAE
Chief, SRAE, NEC

Enrique A. Tayag
Enrique A. Tayag, MD, PHSAE, FPSMID
Director IV, NEC

Philippine HIV & AIDS Registry

The Philippine HIV & AIDS Registry is the official record of the total number of laboratory-confirmed HIV positive individuals, AIDS cases and deaths, and HIV positive blood units in the Philippines. All individuals in the registry are confirmed by the STD/AIDS Cooperative Central Laboratory (SACCL) at San Lazaro Hospital. While all blood units are confirmed by the Research Institute for Tropical Medicine (RITM). Both are National Reference Laboratories (NRL) of the Department of Health (DOH).

Mandatory HIV testing is unlawful in the Philippines (RA 8504). The process of reporting to the Registry is as follows: All blood samples from accredited HIV testing facilities that are screened HIV reactive are sent to SACCL (individuals) or RITM (blood units) for confirmation. Confirmed HIV positive individuals and blood units are reported to the DOH-National Epidemiology Center (NEC), and are recorded in the Registry.

The Registry is a passive surveillance system. Except for HIV confirmation by the NRL, all other data submitted to the Registry are secondary and cannot be verified. An example would be an individual's reported place of residence. The Registry is unable to determine if this reported address is where the person got infected, or where the person lived after being infected, or where the person is presently living, or whether the address is valid. This limitation has major implications to data interpretation. Thus, readers are cautioned to carefully weigh the data and consider other sources of information prior to arriving at conclusions.