
*



HIV/AIDS Situation in Hong Kong [2013]

Annual reported cases reach record high number in 2013

The first case of HIV infection in Hong Kong was reported in 1984. As of 2013, the Department of Health has received a cumulative total of 6,342 reports of HIV infection and 1,437 AIDS cases under the voluntary and anonymous HIV/AIDS reporting system (Fig. 1). The number of annual reports was a record high of 559 cases in 2013, 9% increase compared to 2012. People infected with HIV progress to AIDS when they suffer from clinical complications of severe immunodeficiency due to HIV. In 2013, 84 AIDS reports were received. The most common illnesses presenting at AIDS were *Pneumocystis* pneumonia and tuberculosis.



^{*}AIDS cases are included in the HIV cases

Most infections were in young adult, male and Chinese

A majority of the HIV reports in 2013 are male (79%) and Chinese (70%). Most (84%) infected people were diagnosed at the age between 20 and 49 (Fig.2-4).






Fig 3. HIV reports by ethnicity (2013)
unknown
Chinese
26%
Chinese
70%



Sexual contact remained the major route of transmission for HIV

Sexual contact contributed to around 77% of all reported HIV cases in 2013 (homosexual 49%, heterosexual 25%, bisexual 4%). (Fig 5). There were 7 cases of infection via injecting drug use (IDU) and one case transmitted via blood/blood product infusion outside Hong Kong reported in 2013. 21% of cases in 2013 did not have risk factor ascertained due to inadequate information.


Fig 5. Suspected route of HIV transmission (2013)



Most of the cases were infected locally

In 2013, over half (52%) of reported cases were assessed to have contracted the virus in Hong Kong, 11% in mainland China and in other places respectively. 26% did not have the place of infection ascertained due to inadequate information (Figure 6). In 2013, a majority of MSM cases (74%) contracted the virus in Hong Kong (Fig 7) as compared to less than half (45%) of heterosexual male cases. (Fig 8).

Fig 7. Suspected place of infection MSM (2013)



A continuous rising number of infections among men having sex with men (MSM) is a cause for concern

HIV infection among MSM has shown a continuous rising trend since 2004. Homosexual and bisexual contacts as route of transmission contributed to more than half (53%) of all HIV cases and 66% of the infections in men in 2013 (Fig. 9). The record high of 295 MSM infections reported in 2013 contributed to the high overall number. The number of HIV reports from MSM has been persistently higher than that from heterosexual men since 2005, and the trend continue to widen in the past few years. On the contrast, heterosexual male cases shown a decreasing proportion in past few years (15% in 2013)

Fig 6. Suspected place of infection (2013)




Fig 8. Suspected place of infection Heterosexual male (2013)





Fig 9. HIV reports in heterosexual men and MSM (2004-2013)



In 2013:

- One in every 40,000 new blood donors was tested HIV positive
- One in every 7,000 pregnant women attending public antenatal services was tested HIV positive
- One in every 290 attendees in STD clinics was tested HIV positive
- One in every 240 users in methadone clinics was tested HIV positive

Other publications on HIV/AIDS situation in Hong Kong, such as Annual HIV Surveillance Report, Quarterly STD/AIDS Update are available at http://www.aids.gov.hk

FACTSHEET on HIV/AIDS Situation in Hong Kong [2013]
Prepared by Special Preventive Programme
Centre for Health Protection, Department of Health
Telephone: 3143 7225 Fax: 2780 9580

Website: http://www.aids.gov.hk Email: aids@dh.gov.hk

March 2014