SNAPSHOTS

HIV Epidemic in Asia and the Pacific



How to interact with the data cards




Search and install a QR reader such as QR Reader by Scan

QR Reader by Scan


Scan the QR code using the app

Go to video/webpage link on your device

Asia and the Pacific




Regional overview of trends in HIV infections and AIDS-related deaths




Source: Prepared by www.aidsdatahub.org based on UNAIDS 2016 HIV estimates


Behaviour and response




Proportion of new HIV infections in Asia and the Pacific, 2015



Prevention of mother-to-child transmission (PMTCT)




People on ART: Current pace versus Fast-Track




Source: Prepared by www.aidsdatahub.org based on UNAIDS 2016 HIV estimates and Global AIDS Response Progress Reporting

New HIV infections trend among young people and adults




Less than half of the female sex workers in the region know their HIV status






Consistent condom use is low in many countries




Source: Prepared by www.aidsdatahub.org based on: Integrated Biological and Behavioural Surveys; Behavioural Surveillance Surveys; Li, D., Wang, L., Lin, W., Li, P., Guo, W., et. al; HIV and syphilis infections among street-based female sex workers in China, 2010-2012. Chin Med J 2014; Global AIDS Response Progress Reporting.

Low level of consistent condom use in many countries






Less than half of men who have sex with men know their HIV status





Slow decline in new HIV infections among children





Large gaps in HIV services for mothers and children



40% of HIV+ pregnant women diagnosed



97% of diagnosed HIV+ mothers received ARVs for PMTCT



61% of infants born to diagnosed HIV+ mothers received ARV prophylaxis



48% of infants born to diagnosed HIV+ mothers received HIV test



HIV and Hepatitis C double disease burden





One in three people who inject drugs are in Asia and the Pacific



In Asia and the Pacific...


One in eight people who inject drugs are living with HIV



Among people who inject drugs living with HIV... Nine in ten are co-infected with Hepatitis C




Insufficient coverage of needle and syringe programmes in many countries




Source: Prepared by www.aidsdatahub.org based on: Platt, L., Easterbrook, P., Gower, E., et al. Prevalence and burden of HCV co-infection in people living with HIV: a global systematic review and meta-analysis. *Lancet Infectious Diseases 2016*; Global AIDS Response Progress Report.



Low level of consistent condom use





High levels of violence against transgender people

