

Research article

Open Access

Multilevel analysis of HIV related risk behaviors among heroin users in a low prevalence community

Huizhen Li¹, William Goggins² and Shui Shan Lee*¹

Address: ¹Stanley Ho Centre for Emerging Infectious Diseases, The Chinese University of Hong Kong, Hong Kong, PR China and ²School of Public Health and Nethersole School of Nursing, The Chinese University of Hong Kong, Hong Kong, PR China

Email: Huizhen Li - fallenkerry@hotmail.com; William Goggins - wgoggins@cuhk.edu.hk; Shui Shan Lee* - sslee@cuhk.edu.hk

* Corresponding author

Published: 12 May 2009

Received: 1 September 2008

BMC Public Health 2009, 9:137 doi:10.1186/1471-2458-9-137

Accepted: 12 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2458/9/137>

© 2009 Li et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Injecting drug users (IDU) are at increased risk of human immunodeficiency virus (HIV) infection. Their HIV prevalence however varies from place to place and may not be directly linked with the level of individual risk. This study explores the relative importance of individual and community level characteristics in the practice of HIV-related risk behaviors in IDU in Hong Kong where the HIV prevalence has remained low at below 1%.

Methods: Methadone clinics were used as the channel for accessing drug users in Hong Kong. HIV-related risk factors in drug users attending these clinics were retrieved from a questionnaire routinely administered to newly admitted and readmitted clients, and assessed using logistic regression and multilevel analyses.

Results: Between 1999 and 2005, a total of 41196 person-admissions were recorded by 20 methadone clinics. Male gender, older age and new admissions in bigger clinics located in districts with older median age were more likely to have engaged in HIV related risk behaviors including heroin injection, needle sharing, unprotected sex and having multiple sex partners ($p < 0.05$).

Conclusion: Multilevel analysis is a useful adjunct for determining the association between risk behaviors and both individual and community factors in IDUs, which can be demonstrated even in low HIV prevalence settings.

Background

Injecting drug users (IDUs) are vulnerable to HIV infection both because of the sharing of contaminated injection equipment and their practice of high risk sexual behaviors. In Hong Kong, a metropolitan city in Southern China, only 5.6% of the 3400 HIV cases reported as of mid-2007 were IDUs, and their HIV prevalence has remained at no more than 1% in different settings.[1] However, some Chinese provinces have reported HIV prevalences of more than 20% in IDUs, and in most cities

in the Pearl River Delta region adjacent to Hong Kong the estimated prevalences have ranged from 3–6%. [2-4] HIV outbreaks can potentially occur in future because some IDUs may have risky practices in China. A series of local studies conducted between 2000 and 2004 revealed that 25% of Hong Kong's IDUs had ever used drugs in Mainland China.[5] Other reports suggested that Mainland Chinese commercial sex workers (CSWs) who were also IDUs were less inclined to use condoms (50% vs 77%) and more likely to share needles (50% vs 20%), compared

with non drug using CSWs.[4,6] Chances are that if IDUs in Hong Kong inject and share needles with Mainland Chinese IDUs in high HIV prevalence areas in China, or have unprotected sex with CSWs, HIV could easily spread. An assessment of the risk behaviors of IDUs in Hong Kong is important for predicting the potential of HIV spread, despite the low prevalence currently recorded locally.

IDUs constitute a marginalized and hard-to-reach population in society. They tend to socialize with their fellow IDUs rather than members of the general population. This unique social network configuration may shape their behavior patterns. Within the same group of IDUs, behaviors tend to be similar because of peer influence. The extent to which IDUs engage in risky behaviors (injection, needle sharing, syringe reuse, multiple sex partners and unprotected sex), and their individual demographic characteristics, such as age, gender, ethnicity, have been found to be correlated with HIV seropositivity. [7-9] A study conducted in the United States [10] considered IDUs with similar practices as members of a 'homogeneous cluster', and went on to describe the variation of HIV seroprevalence across clusters. Another study [11] reported that community characteristics derived from census data were predictors of HIV related risk among people residing in the same census tract. These multilevel studies were conducted in western countries, and so far no similar findings have been reported from Asian countries. In this study, we undertook to examine the variation of IDU associated risk behaviors at individual and community level in an Asian city, Hong Kong. Methadone clinics were used as the channel for accessing IDUs because of their high coverage of heroin users in the territory. The government-run clinics were set up in 1970s, and over the past three decades they have evolved to become focal points of neighborhoods for IDUs in different geographic areas. Service data available from these clinics were used to assess the relative importance of individual and community level characteristics on the practice of risk behaviors.

Methods

This was a multi-site, institution based, cross-sectional study using data retrieved from results of a routine structured questionnaire administered on clients of the 20 methadone clinics in Hong Kong following informed consent. Institutional approval for access to the data was sought from the Department of Health, in compliance with the Personal Data (Privacy) Ordinance. The conduct of the study complied with the provisions of the Declaration of Helsinki. Contents of the questionnaire included anonymized personal data, and questions on drug taking habits, and sexual practices in the 12-month period prior to registration. There were two categories of registration in methadone clinics, new admission and readmission. New admissions referred to those clients who have never taken methadone treatment before, whereas readmissions were

those who had enrolled in methadone treatment in the past, but had dropped out for over 28 days. Since each entry was anonymous, we could not match the readmissions with the new admissions of the same individuals. Therefore, in this study, only data from questionnaires on newly admitted drug users were used in logistic regression and multilevel analyses.

Risk behaviors were assessed to determine the vulnerability of drug users to HIV infection. Although early studies have determined the relative transmission efficiencies of HIV for each practice, [12-14] recent studies demonstrated the importance of sexual contact as a risk factor as compared to injection in IDUs.[15] Practice of injection, needle sharing, unprotected sex and multiple sex partners were assessed individually. Injection was defined as any report of injecting drug prior to registration, while needle sharing behavior was divided into current and past sharing practice. Unprotected sex referred to the use of condoms in less than 50% of the sexual activities in the preceding 12 months, while those reporting to have more than 1 sex partner in the same time period was defined as having multiple sex partners.

Two sets of variables were computed to determine their roles as predictors – at individual level and community level. Taking reference from variables that had been found to be significant in other studies, the following individual level predictors were used: gender, age and admission quarter/year. Age and its square were used to explore the non-linear association of age with HIV risk behaviors. Community level variables included population level data from each of the 18 administrative districts in Hong Kong obtained from the 2006 by-census. Only median age, median household income level, population density, and education level were available from this database. As income was highly correlated with education level, only income level was used in the analysis. In addition, the number of total admissions for each clinic was used as a community level variable reflecting the size of the IDU neighborhood.

Logistic regression was used to examine the effects of individual level predictor variables on the binary outcomes. Significance levels were reported for multivariate analyses and multilevel analyses with statistical significance set at 0.05. Age and age-squared were entered into the models as continuous variables and odds ratios were calculated for selected ages with age 20 as the reference. SPSS version 13.0 (SPSS Inc 2004) was used for the descriptive statistical analyses and to fit the logistic regression models. Multilevel logistic regression models were used to assess the independent predictive ability of both individual and community level covariates, and were fit using Supermix 1 statistical software (SSI Inc).

Results

Additional file 1 provides a descriptive overview of the characteristics of our study population. From 1999 to 2005, there were totally 41196 person-admissions, of which 5184 were newly admitted drug users. The new admissions were predominantly male (77.6%) and aged below 30 (62.4%). They were younger than drug users on readmission (in which 53.7% were aged between 30 and 50). The yearly number of new admissions dropped steadily from around 1004 in 1999 to 482 in 2005, with a rebound to 965 in 2002.

The most frequently reported drug of abuse was heroin. More than 98% of the new admission clients reported the use of heroin alone or in combination with other drugs. Their commonest HIV related risk behaviors were injection and unprotected sex. While a majority of the new admissions preferred inhalation, smoking or sniffing, more than half of the readmission cases used injection as their major mode of heroin use. A quarter of the newly admitted heroin users reported current injection. Unclean syringe use was reported by 18% of new admissions, while the proportion among readmissions was 14%. Current needle sharing was reported in only 2% of the new admissions and readmissions, but some 7% of the readmissions reported past needle sharing. A slightly higher proportion of new admissions than readmissions reported having multiple sex partners while the reversed was true for unprotected sex in the preceding 12 months (10.1% vs 8.8%, $p > 0.05$ for multiple sex partners and 61.6% vs 63.7%, $p = 0.02$, odds ratio (OR): 0.91, 95% confidence interval (CI): 0.84–0.98 for unprotected sex).

Using multivariable logistic regression, 4 separate models – with injection, needle sharing, unprotected sex and multiple sex partners as outcomes – were used to assess HIV related risks. (Additional file 2) Male gender was strongly associated with heroin injection and having multiple sex partner (adjusted odds ratio (AOR): 1.27, 95%CI: 1.08–1.48, $p = 0.004$; AOR: 2.54, 95%CI: 1.91–3.38, $p < 0.001$ respectively) in all except the needle sharing model, in which only male gender was correlated with needle sharing behavior in univariate analysis. However, male drug users were less likely to have engaged in unprotected sex than female in the study population (AOR: 0.60, 95%CI: 0.52–0.68, $p < 0.001$). Age was also a significant predictor of heroin injection and unprotected sex. The positive coefficient for the age term and the negative coefficient for the age-squared term suggested a non-linear association, with more risky behaviors occurring in the middle of the age ranges (30 to 40 for injecting heroin use and 30 to 50 for unprotected sex). The time trend was evaluated by assessing the role of admission date. Later admission quarter/year was associated with more injection and lower tendency of having multiple sex partners ($p < 0.001$ in both models).

Preliminary analyses were performed on the data prior to multilevel modeling, the results of which confirmed the contextual structure of data and demonstrated the existence of between-community variation. In applying the multilevel model, age, gender, and admission were considered to exert fixed effects. Community-level variables (household income, clinic size, population density and median age of district population) were included as predictors of district-specific intercept, while no variables were used to predict the slope for time because no significant cross-level interactions were found in the preliminary analysis. Results of our models (Additional file 3) showed that individual variables (gender, age and admission time) remained as significant factors for predicting the risk behavior. Male gender was significantly associated with all four HIV risk behaviors, while the significance of age was seen only in the injection and unprotected sex models. Later admission was associated with slightly more injection practice and lower likelihood of multiple sex partners. After controlling for individual characteristics including age and gender, clients from bigger clinics, as defined by high caseload, were more likely to have injected heroin, shared needle and had sex without condom ($p < 0.05$). Clients from districts with older median age were more likely to have practiced risky sexual behaviors ($p < 0.001$).

Discussion

This is the first study in an Asian population involving the analysis of data from methadone clinics to examine the variation of risk behaviors of drug users, using both logistic regression models and multilevel models. Results from our study demonstrated a variation of HIV risk behaviors across neighborhoods, as defined by the clinic locations where methadone users went for regular treatment. In Hong Kong, each methadone clinic functions like the center of a separate neighborhood. Clients of most clinics resided near the clinics at which they registered. Apparently, drug users spent most of their time around the clinic area and socialized with other drug users from the same neighborhood. It is not surprising, therefore, that they formed a local social network and were under the influence of their fellow drug users, leading to a similar pattern of drug taking behaviors. Associated with drug injection, needle sharing and unprotected sex, bigger clinic size was a predictive factor of risk behavior in our multilevel models. This result was consistent with studies elsewhere,[16,17] in which smaller social network implied lower levels of risky drug use behaviors. We did not, however, have relevant data for assessing the exact social network size of each drug user, and the inter-relationship between network size and with clinic size. The possible association of clinic size with HIV risk would need to be investigated in a study designed specifically for this purpose. As a harm reduction measure, the optimum size of




Figure 1
Geographic distribution of methadone clinics in Hong Kong.

methadone clinics is important, in order that maximum benefits of substitution treatment can be derived without engendering undesirable behavioral risks.

Some studies had examined the association between population characteristics and injection practice among IDUs.[11,18] It was shown in our study that an older median age in a district was a predictor of sexual risk. Median population age reflects the composition of population in the locale, as well as the education level and income level. People with older age were less likely to have been exposed to safer sex messages which had become standard publicity themes only after the beginning of the AIDS epidemic in early 80s. It may therefore be argued that condom use and multiple sex partners could in fact be more common in the older population, suggesting that drug users in such neighborhoods could be influenced to share the same practices. Furthermore,

median population age could also be a surrogate for population density, since in Hong Kong the districts with older populations tend to be the more densely populated older parts of the city. In our study however population density was not found to be a significant predictor. Using multilevel analysis, we failed to find any association between population income and HIV risk. In our study, median household income was used as the surrogate, the level of which ranging between HKD9000 and HKD30000 (USD1 = HKD7.8). Since individual income level was not available and thus not controlled in the multilevel model, the analysis might not be robust enough to detect any effect of poverty on risk behaviors. Neighborhood variation in HIV prevalence has in fact been demonstrated in a previous study.[19] Among the neighborhood conditions, lower income was found to be associated with higher HIV prevalence among IDUs.[20] Overall, we were able to demonstrate a variation of HIV risk at community

level. The generally low HIV rate in drug users in Hong Kong does not allow us to examine the correlation of such factors with HIV prevalence.

On an individual level, our study showed a strong association of male gender with heroin injection, needle sharing and higher sexual risk, while age (30 to 40 for drug injection and older age for unprotected sex) also exhibited a positive association. Male gender and older age have been shown to be predictive factors of increased HIV risk in other studies.[21,22] A similar study using existing record to explore the behavioral change of IDUs in Hong Kong demonstrated a decline in needle sharing over the years,[23] an observation also made in our study. The low prevalence of needle sharing in our study population may be a result of the easy access to disposable syringes from local pharmacies. Also, the good coverage of methadone services has helped to reduce needle sharing behaviors.(Figure 1) Taken together, drug users' access to syringe and the provision of substitution treatment at the methadone services provided an explanation for the low HIV rate in Hong Kong.[24]

Our multilevel study of drug users carried certain limitations. First of all, Hong Kong is a small city with a total area of just 1000 Km² and with relatively 'homogenous' community characteristics. Unlike countries such as the United States in which the IDU population is often composed of a variety of ethnicities, almost all drug users in Hong Kong are Chinese. Variation of ethnicities has been reported to be related to HIV transmission in IDU,[25] a phenomenon that should be less important in Hong Kong. Our analysis illustrates community level variation in risk behaviors between clinics despite the 'homogenous' community characteristics and thus provides evidence that multilevel modeling could also be applied to community with less variability of the population profile. On the other hand, the study sample might not be representative of all drug users in the territory. Theoretically a representative sample could have been obtained by conducting street surveys in randomly selected sites in the city. [26-28] The feasibility of such sampling is low because IDUs constitute a hard-to-reach population, and social discrimination often creates an unfriendly environment for them to be accessed. Furthermore, their use of illicit drugs dissuades them to reveal their lifestyle to strangers.[29] Our solution to the sampling problem was to resort to existing service data.[30] In this connection, the government-run methadone treatment service in Hong Kong provides a particularly useful data source. The clinics have been in contact with an estimated 70% of all drug users.[31] Each methadone user is interviewed on admission and their demographic data, drug taking habits and sexual behaviors are recorded. With different opening hours, the 20 methadone clinics are widely distributed

and are convenient for both jobless and employed clients. The major limitation is however the availability of small number of selected predefined fields for analyses. Another possible limitation is the self-reported nature of the data. Although it is mandatory for the questionnaire to be completed on admission, the self-reported behaviors may not be entirely reliable. As methadone users have to register their names and identity card numbers (anonymized in this study), there is the possibility that they are inclined to conceal their practices, though the prescription of methadone doses is not linked with reported behaviors. An underestimation of practice of risky behaviors is nevertheless a cause for concern. In some studies, however, self-reports of behaviors were found to be reliable in the collection of data on risk behaviors.[32,33]

Conclusion

The increasing reports of HIV infections through injecting drug use among the HIV cases in Hong Kong raises the concern about the potential spread of the virus within this population and beyond.[34] Results from our study have provided an analytical description of the characteristics of the HIV related risk behaviors of IDUs. By elucidating the association between risk behaviors and both individual and community level factors, our approach may help predict the HIV risk of IDUs in a spatial context, which can in turn support effective allocation of resources to achieve prevention. We have demonstrated that the application of multilevel modeling could explain the contextual influences of individual and community factors on HIV related behavior in an Asian population, thereby calling for interventions to be geared towards this new angle in order to achieve effective prevention.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HL conducted field study, collated the data, conducted all statistical analyses, and prepared the first draft of the manuscript. WG provided statistical support to the study. SSL conceptualized the study, coordinated all research activities and edited the manuscript. All authors have read the revised version of the manuscript and approved the final version.

Additional material

Additional File 1

Characteristics of the study population. The data provided shows the demographics, drug taking habits, and the practice of high risk behaviors in drug users in the study population.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2458-9-137-S1.doc>]

Additional File 2

Predictors of HIV risk at individual level. Results of logistic regression with OR (95%CI) for determining possible predictors of HIV risk.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2458-9-137-S2.doc>]

Additional File 3

Predictors of HIV risk at two levels. Results of multilevel analysis with OR (95%CI) (N = 5160) for determining possible predictors for HIV risk.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2458-9-137-S3.doc>]

Acknowledgements

The authors would like to thank Department of Health, Hong Kong SAR Government, for access to the anonymised methadone clinics data, Mr. Liu at the Society for the Aid and Rehabilitation of Drug Abusers for the arrangement of volunteer helpers from Project Phoenix to advise on the interpretation of research results, SpatioEpi Group for preparing the map, and staff at Stanley Ho Centre for Emerging Infectious Diseases, The Chinese University of Hong Kong, for their administrative support in the course of conducting the study. Contents of this report also contributed partly to the Master of Philosophy thesis of Huizhen Li at The Chinese University of Hong Kong.

References

- Department of Health Hong Kong: *HIV Surveillance Report – 2005 update. Hong Kong 2006* [<http://www.info.gov.hk/aids/archives/backissues/taids05.pdf>].
- State Council AIDS Working Committee Office and UN Theme Group on HIV/AIDS in China: *A Joint Assessment of HIV/AIDS Prevention, Treatment and Care in China* Beijing: State Council AIDS Working Committee Office; 2004.
- Mingjian N, Wheeler KM, Cheng J, Yonghai D, Chen W, Fitzwarryne C, Wang J: **HIV/AIDS prevalence and behaviour in drug users and pregnant women in Kashgar Prefecture: case report.** *Harm Reduct J* 2006, **3**:7.
- Monitoring the AIDS Pandemic Network: *MAP Report 2005: Drug Injection and HIV/AIDS in Asia 2005* [http://www.mapnetwork.org/docs/MAP_IDU%20Book%2024Jun05_en.pdf].
- Working Group on HIV Prevention for Commercial Sex Workers and their Clients, Community Forum on AIDS, and Hong Kong Advisory Council on AIDS: *Report of Community Assessment and Evaluation of HIV Prevention for Commercial Sex Workers and their Clients in Hong Kong 2006. Hong Kong 2006* [<http://www.info.gov.hk/aids/pdf/g169.pdf>].
- Monitoring the AIDS Pandemic Network: *MAP Report: Sex work and HIV/AIDS in Asia 2005* [http://www.mapnetwork.org/docs/MAP_SVW%20in%20Asia%20Final%2004July05_en.pdf].
- Perngmark P, Celentano DD, Kawichai S: **Risk factors for HIV infection among drug injectors in southern Thailand.** *Drug Alcohol Depend* 2003, **71**:229-238.
- Liu W, Chen J, Rodolph M, Beauchamp G, Masse B, Wang S, Li R, Ruan Y, Zhou F, Leung MK, Lai S, Shao Y, Jackson JB: **HIV prevalence among injection drug users in rural Guangxi China.** *Addiction* 2006, **101**:1493-1498.
- Hien NT, Giang LT, Binh PN, Deville W, van Ameijden EJ, Wolffers I: **Risk factors of HIV infection and needle sharing among injecting drug users in Ho Chi Minh City, Vietnam.** *J Subst Abuse* 2001, **13**:45-58.
- Williams ML, Zhao Z, Freeman RC, Elwood WN, Rusek R, Booth RE, Dennis ML, Fisher DG, Rhodes F, Weatherby NL: **A cluster analysis of not-in-treatment drug users at risk for HIV infection.** *Am J Drug Alcohol Abuse* 1998, **24**:199-223.
- Bluthenthal RN, Do DP, Finch B, Martinez A, Edlin BR, Kral AH: **Community characteristics associated with HIV risk among injection drug users in the San Francisco Bay Area: a multi-level analysis.** *J Urban Health* 2007, **84**:653-666.
- Kaplan EH, Heimer R: **HIV incidence among New Haven needle exchange participants: updated estimates from syringe tracking and testing data.** *J Acquir Immune Defic Syndr Hum Retroviral* 1995, **10**:175-176.
- UNAIDS: *AIDS epidemic update: special report on HIV/AIDS: December 2006 2009* [<http://www.unaids.org/en/KnowledgeCentre/HIVData/EpiUpdate/EpiUpdArchive/2006/default.asp>]. Geneva: UNAIDS and WHO
- Varghese B, Maher JE, Peterman TA, Branson BM, Steketee RW: **Reducing the risk of sexual HIV transmission: quantifying the per-act risk for HIV on the basis of choice of partner, sex act, and condom use.** *Sex Transm Dis* 2002, **29**:38-43.
- Kral AH, Bluthenthal RN, Lorvick J, Gee L, Bacchetti P, Edlin BR: **Sexual transmission of HIV-1 among injection drug users in San Francisco, USA: risk-factor analysis.** *Lancet* 2001, **357**:1397-1401.
- Suh T, Mandell W, Latkin C, Kim J: **Social network characteristics and injecting HIV-risk behaviors among street injection drug users.** *Drug Alcohol Depend* 1997, **47**:137-143.
- Zapka JG, Stoddard AM, McCusker J: **Social network, support and influence: relationships with drug use and protective AIDS behavior.** *AIDS Educ Prev* 1993, **5**:352-366.
- Wang J, Siegal HA, Falck RS, Carlson RG: **Needle transfer among injection drug users: a multilevel analysis.** *Am J Drug Alcohol Abuse* 1998, **24**:225-237.
- Poundstone KE, Strathdee SA, Celentano DD: **The social epidemiology of human immunodeficiency virus/acquired immunodeficiency syndrome.** *Epidemiol Rev* 2004, **26**:22-35.
- Friedman SR, Lieb S, Tempalski B, Cooper H, Keem M, Friedman R, Flom PL: **HIV among injection drug users in large US metropolitan areas, 1998.** *J Urban Health* 2005, **82**:434-445.
- Zhang Y, Shan H, Trizzino J, Ruan Y, Beauchamp G, Masse B, Ma J, Gu Y, He Y, Rui B, Wang J, Poundstone K, Jiang Y, Brooks JJ, Shao Y: **Demographic characteristics and risk behaviors associated with HIV positive injecting drug users in Xinjiang, China.** *J Infect* 2007, **54**:285-290.
- Williams ML, McCurdy SA, Atkinson JS, Kilonzo GP, Leshabari MT, Ross MW: **Differences in HIV risk behaviors by gender in a sample of Tanzanian injection drug users.** *AIDS Behav* 2007, **11**:137-144.
- Lee SS, Hollinrake JB, NG MH: **Changing behavioural pattern of drug users in Hong Kong 1991-1995.** *Addiction* 1998, **93**:541-548.
- Van Den BC, Smit C, Van BG, Coutinho R, Prins M: **Full participation in harm reduction programmes is associated with decreased risk for human immunodeficiency virus and hepatitis C virus: evidence from the Amsterdam Cohort Studies among drug users.** *Addiction* 2007, **102**:1454-1462.
- Centers for Disease Control and Prevention: *HIV/AIDS Surveillance Report, 2005. Atlanta 2007* [<http://www.cdc.gov/hiv/topics/surveillance/resources/reports/2005report/pdf/2005SurveillanceReport.pdf>].
- Hunter GM, Stimson GV, Judd A, Jones S, Hickman M: **Measuring injecting risk behaviour in the second decade of harm reduction: a survey of injecting drug users in England.** *Addiction* 2000, **95**:1351-1361.
- Perngmark P, Celentano DD, Kawichai S: **Needle sharing among southern Thai drug injectors.** *Addiction* 2003, **98**:1153-1161.
- Peters A, Davies T, Richardson A: **Multi-site samples of injecting drug users in Edinburgh: prevalence and correlates of risky injecting practices.** *Addiction* 1998, **93**:253-267.
- Bobrova N, Rhodes T, Power R, Alcorn R, Neifeld E, Krasniukov N, Latyshevskaya N, Maksimova S: **Barriers to accessing drug treatment in Russia: a qualitative study among injecting drug users in two cities.** *Drug Alcohol Depend* 2006, **82**(Suppl 1):S57-S63.
- Archibald CP, Jayaraman GC, Major C, Patrick DM, Houston SM, Sutherland D: **Estimating the size of hard-to-reach popula-**

tions: a novel method using HIV testing data compared to other methods. *AIDS* 2001, **15(Suppl 3)**:S41-S48.

31. Narcotics Division Security Bureau HKSAR: *Central Registry of Drug Abuse Fifty-fifth Report*. Hong Kong 2006 [http://www.nd.gov.hk/55_report.htm].
32. Edwards JW, Fisher DG, Johnson ME, Reynolds GL, Redpath DP: **Test-retest reliability of self-reported drug treatment variables**. *J Subst Abuse Treat* 2007, **33**:7-11.
33. Schrimshaw EV, Rosario M, Meyer-Bahlburg HF, Scharf-Matlick AA: **Test-retest reliability of self-reported sexual behavior, sexual orientation, and psychosexual milestones among gay, lesbian, and bisexual youths**. *Arch Sex Behav* 2006, **35**:225-234.
34. Department of Health Hong Kong: *Hong Kong STD/AIDS update, a quarterly surveillance report: Quarter 4 2007*. Hong Kong 2007, **13(4)**: [<http://www.info.gov.hk/aids/archives/backissuestd/std07q4.pdf>].

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/9/137/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

