

Survey: National Baseline High Risk Groups and Bridge Population BSS, India 2002: Men Who Have Sex with Men

Title:	Men Who Have Sex with Men													
Survey Type:	BSS													
Country:	India													
Year:	2002													
Start Date:	February 2002													
End Date:	March 2002													
Respondents:	<table border="1"> <thead> <tr> <th>Female</th> <th>Male</th> </tr> </thead> <tbody> <tr> <td>Respondent: [n/a]</td> <td>Respondent: Target Groups</td> </tr> <tr> <td>Age: [n/a]</td> <td>Age: 12 - 60</td> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: 1387</td> </tr> <tr> <th>Facilities</th> <th>Households</th> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: [n/a]</td> </tr> </tbody> </table>	Female	Male	Respondent: [n/a]	Respondent: Target Groups	Age: [n/a]	Age: 12 - 60	Sample Size: [n/a]	Sample Size: 1387	Facilities	Households	Sample Size: [n/a]	Sample Size: [n/a]	
Female	Male													
Respondent: [n/a]	Respondent: Target Groups													
Age: [n/a]	Age: 12 - 60													
Sample Size: [n/a]	Sample Size: 1387													
Facilities	Households													
Sample Size: [n/a]	Sample Size: [n/a]													
Implementing:	National AIDS Control Organization, ORG-CSR, India													
Description:	National Baseline BSS among Men Who Have sex with Men(MSM) was conducted in 5 cities/States across the country: Delhi, Mumbai, Kolkatta, Chennai, Bangalore. Topics included in the questionnaire: Background Characteristics, Knowledge, Opinion and Attitude towards STI/HIV/AIDS, Sexual History, Male Partner, Risk Perception and Exposure to Intervention.													
Footnote:	[n/a]													

Indicators:	<p>4.0.1) Heard of HIV/AIDS (97%)</p> <p>4.1) Knowledge of HIV prevention methods - Composite of 2 components (prompted) (69%)</p> <p>4.1.2) Knowledge of HIV prevention methods - Use of condoms (prompted) 83%</p> <p>4.1.3) Knowledge of HIV prevention methods - Only one partner (prompted) 73%</p> <p>4.1.4) Knowledge of HIV prevention methods - Abstain from sex (prompted) 74%</p> <p>4.2) No incorrect beliefs about AIDS - Composite of 3 components 54%</p> <p>4.2.1) No incorrect beliefs about AIDS - Healthy-looking person can have the AIDS virus 71%</p> <p>4.2.2) No incorrect beliefs about AIDS - AIDS cannot be transmitted by mosquito bites 78%</p> <p>4.2.6) No incorrect beliefs about AIDS - Cannot become infected by sharing food with someone who has AIDS (84%)</p> <p>4.5.4) Knowledge of MTCT - During pregnancy (89%)</p> <p>4.5.6) Knowledge of MTCT - Through breastfeeding (72%)</p> <p>5.1) Population requesting an HIV test, receiving a test and receiving test results (30%)</p> <p>5.1.1) Population receiving an HIV test (35%)</p> <p>8.1) Higher risk sex in the last year (79%)</p> <p>8.2.2) Condom use at last sex with anyone (36%)</p> <p>8.3) Commercial sex in last year (36%)</p>
--------------------	--

8.4)	Condom use at last commercial sex, reported by client 39%
8.7)	Condom use at last anal sex between men 53%
12.4)	Men and women seeking treatment for STIs 72%

Survey: National Baseline High Risk Groups and Bridge Population BSS, India 2002: Injecting Drug Users (IDUs)

Title:	Injecting Drug Users (IDUs)												
Survey Type:	BSS												
Country:	India												
Year:	2002												
Start Date:	February 2002												
End Date:	March 2002												
Respondents:	<table border="1"> <thead> <tr> <th>Female</th> <th>Male</th> </tr> </thead> <tbody> <tr> <td>Respondent: [n/a]</td> <td>Respondent: Target Groups</td> </tr> <tr> <td>Age: [n/a]</td> <td>Age: 12 - 60</td> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: 1355</td> </tr> <tr> <th>Facilities</th> <th>Households</th> </tr> <tr> <td>Sample Size: [n/a]</td> <td>Sample Size: [n/a]</td> </tr> </tbody> </table>	Female	Male	Respondent: [n/a]	Respondent: Target Groups	Age: [n/a]	Age: 12 - 60	Sample Size: [n/a]	Sample Size: 1355	Facilities	Households	Sample Size: [n/a]	Sample Size: [n/a]
Female	Male												
Respondent: [n/a]	Respondent: Target Groups												
Age: [n/a]	Age: 12 - 60												
Sample Size: [n/a]	Sample Size: 1355												
Facilities	Households												
Sample Size: [n/a]	Sample Size: [n/a]												
Implementing:	National AIDS Control Organization, India												
Description:	National Baseline BSS among Men Injecting Drug Users(IDU) was conducted in 5 cities/States across the country:Delhi, Mumbai, Kolkata, Chennai, Manipur. Topics included in the questionnaire: Background Characteristics, Drug Use, Needle and Syringe Sharing Behaviour, Knowledge, Opinion and Attitude towards STI/HIV/AIDS, Sexual History, Risk Perception and Exposure to Intervention.												
Footnote:	[n/a]												
Indicators:	<p>4.0.1) Heard of HIV/AIDS 97%</p> <p>4.1) Knowledge of HIV prevention methods - Composite of 2 components (prompted) 65%</p> <p>4.1.2) Knowledge of HIV prevention methods - Use of condoms (prompted) 84%</p> <p>4.1.3) Knowledge of HIV prevention methods - Only one partner (prompted) 71%</p> <p>4.1.4) Knowledge of HIV prevention methods - Abstain from sex (prompted) 65%</p> <p>4.2) No incorrect beliefs about AIDS - Composite of 3 components 43%</p> <p>4.2.1) No incorrect beliefs about AIDS - Healthy-looking person can have the AIDS virus 71%</p> <p>4.2.2) No incorrect beliefs about AIDS - AIDS cannot be transmitted by mosquito bites 67%</p> <p>4.2.6) No incorrect beliefs about AIDS - Cannot become infected by sharing food with someone who has AIDS 69%</p> <p>4.4.1) Knowledge of HIV prevention among injecting drug users - switching to non-injectable drugs 55%</p> <p>4.5.6) Knowledge of MTCT - Through breastfeeding 74%</p>												

- 5.1) [Population requesting an HIV test, receiving a test and receiving test results](#) 20%
- 5.1.1) [Population receiving an HIV test](#) 26%
- 8.1) [Higher risk sex in the last year](#) 10%
- 8.2) [Condom use at last higher risk sex](#) 34%
- 8.2.1) [Condom use at last sex with a spouse or cohabiting partner](#) 34%
- 8.3) [Commercial sex in last year](#) 16%, n=1355
- 8.4) [Condom use at last commercial sex, reported by client](#) 60%, n=217
- 10.1) [Injecting drug users sharing equipment at last injection](#) 41%
- 10.2) [Injecting drug users never sharing equipment in the last month](#) 44%
- 12.4) [Men and women seeking treatment for STIs](#) 35%, n=350